



This PDF file is an excerpt from the EPA report entitled *Wadeable Streams Assessment: A Collaborative Survey of the Nation's Streams* EPA 841-B-06-002 (April 2006). The entire document can be downloaded from <http://www.epa.gov/owow/streamsurvey/report.pdf>.

Wadeable Streams Assessment: A Collaborative Survey of the Nation's Streams EPA 841-B-06-002

Chapter 5 Sources and References

May 2006

Chapter 5 – Sources and References

General References

Griffith, G., Loveland, T., Olsen, T., Omernik, J., (contributors). 1997. *Ecological Regions of North America – Toward a Common Perspective*. North American Commission for Environmental Cooperation.

National Geographic Society. 1988. Close-Up: U.S.A. (maps).

North American Ecological Regions GIS map files - North American Commission for Environmental Cooperation.

Paul, RK and RA Paul. 1977. Geology of Wisconsin and Upper Michigan. Kendall/Hunt Publishing Company.

Population estimates based on U.S. Census Bureau Census 2000 SF 1 Tables for Total Population by County.

Stoddard, J.L., Kahl, J.S., Deviney, F.A., DeWalle, D., Driscoll, C.T., Herlihy, A., Kellogg, J.H., Murdoch, P., Webb, J.R., and Webster, K. 2003. "Response of surface water chemistry to the Clean Air Act Amendments of 1990." U.S. Environmental Protection Agency, Washington, D.C. 79 pp. EPA/620/R-03/001

U.S. Census Bureau TIGER line maps: 1990 U.S. Counties.

Wohl, Ellen E., 2004. *Disconnected Rivers – Linking Rivers to Landscapes*. Yale University.

EMAP Stream and River Sampling Methods

Peck, D. V., Averill, D. K., Herlihy, A. T., Hughes, R. M., Kaufmann, P. R., Klemm, D. J., Lazorchak, J. M., McCormick, F. H., Peterson, S. A., Cappaert, M. R., Magee, T. & Monaco, P. A. (2005). Environmental Monitoring and Assessment Program - Surface Waters Western Pilot Study: Field Operations Manual for Non-Wadeable Rivers and Streams. EPA Report EPA 600/R-05/xxx, U.S. Environmental Protection Agency, Washington, DC.

Peck, D. V., Herlihy, A. T., Hill, B. H., Hughes, R. M., Kaufmann, P. R., Klemm, D. J., Lazorchak, J. M., McCormick, F. H., Peterson, S. A., Ringold, P. L., Magee, T. & Cappaert, M. R. (2005). Environmental Monitoring and Assessment Program - Surface Waters Western Pilot Study: Field Operations Manual for Wadeable Streams. EPA Report EPA 600/R-05/xxx, U.S. Environmental Protection Agency, Office of Research and Development, Washington, DC.

Probability Designs

- Olsen, A. R., Sedransk, J., Edwards, D., Gotway, C. A., Liggett, W., Rathbun, S., Reckhow, K. H. & Young, L. J. (1999). Statistical issues for monitoring ecological and natural resources in the United States. *Environmental Monitoring and Assessment* 54, 1-45.
- Stevens Jr., D. L. (1997). Variable density grid-based sampling designs for continuous spatial populations. *Environmetrics* 8, 167-195.
- Stevens Jr., D. L. & Urquhart, N. S. (2000). Response designs and support regions in sampling continuous domains. *Environmetrics* 11, 11-41.

Ecological Regions

- Omernik, J. M. (1987). Ecoregions of the conterminous United States. *Annals of the Association of American Geographers* 77, 118-125.

Indices of Biotic Integrity

- Barbour, M. T., Stribling, J. B. & Karr, J. R. (1995). Multimetric approach for establishing biocriteria and measuring biological condition. In *Biological assessment and criteria: tools for water resource planning and decision making*. (Davis, W. S. & Simon, T. P., Eds), pp. Chapter 6, pg. 63-77. Lewis, Boca Raton, FL.
- Frey, D. G. (1977). The integrity of water - an historical approach. In *The Integrity of Water*. (Ballantine, S. K. & Guarala, L. J., Eds), pp. 127-140. U.S. Environmental Protection Agency, Washington DC.

- Karr, J. R. & Dudley, D. R. (1981). Ecological perspective on water quality goals. *Environmental Management* 5, 55-68.

- Karr, J. R. (1981). Assessment of biotic integrity using fish communities. *Fisheries* 6, 21-27.

Observed/Expected Models

- Hawkins, C. P. (In Press (2005)). Quantifying biological integrity with predictive models: comparisons with three other assessment methods. *Ecological Applications*.

- Hawkins, C. P., Norris, R. H., Hogue, J. N. & Feminella, J. W. (2000). Development and evaluation of predictive models for measuring the biological integrity of streams. *Ecological Applications* 10, 1456-1477.

- Van Sickle, J., Hawkins, C. P., Larsen, D. P. & Herlihy, A. T. (2005). A null model for the expected macroinvertebrate assemblage in streams. *Journal of the North American Benthological Society* 24, 178-191.

Wright, J. F. (2000). An introduction to RIVPACS. In Assessing the Biological Quality of Fresh Waters. (Wright, J. F., Sutcliffe, D. W. & Furse, M. T., Eds), pp. 1-24. Freshwater Biological Association, Ambleside, UK.

Physical Habitat

Kaufmann, P. R., Levine, P., Robison, E. G., Seeliger, C. & Peck, D. (1999). Quantifying Physical Habitat in Wadeable Streams. EPA Report EPA/600/3-88/021a, U.S. EPA, Washington, D.C.

Reference Condition

Bailey, R. C., Norris, R. H. & Reynoldson, T. B. (2004). Bioassessment of Freshwater Ecosystems: Using the Reference Condition Approach. Kluwer Academic Publishers, New York.

Hughes, R. M. (1995). Defining acceptable biological status by comparing with reference conditions. In Biological Assessment and Criteria: Tools for Water Resource Planning and Decision Making for Rivers and Streams. (Davis, W. & Simon, T., Eds), pp. Chapter 4, pg. 31-47. Lewis, Boca Raton, FL.

Stoddard, J. L., Larsen, D. P., Hawkins, C. P., Johnson, R. K. & Norris, R. H. (In Press (2005)). Setting expectations for the ecological condition of running waters: the concept of reference condition. Ecological Applications.

Other EMAP Assessments

Stoddard, J. L., Herlihy, A. T., Hill, B. H., Hughes, R. M., Kaufmann, P. R., Klemm, D. J., Lazorchak, J. M., McCormick, F. H., Peck, D. V., Paulsen, S. G., Olsen, A. R., Larsen, D. P., Van Sickle, J. & Whittier, T. R. (In Press). Mid-Atlantic Integrated Assessment (MAIA)--State of the Flowing Waters Report. EPA Report EPA 600/R-05/xxx, U.S. Environmental Protection Agency, Washington, DC.

U.S. Environmental Protection Agency (2000). Mid-Atlantic Highlands Streams Assessment. p. 64. EPA Report EPA/903/R-00/015, U.S. Environmental Protection Agency, Region 3, Philadelphia, PA.

Biological Condition Gradient/Quality of Reference Sites

Davies, S. P. & Jackson, S. K. (In press). The Biological Condition Gradient: A conceptual model for interpreting detrimental change in aquatic ecosystems. Ecological Applications.

Lattin, P. D. (In Preparation). A process for characterizing watershed level disturbance using orthophotos.

Relative Risk

Van Sickle, J., Stoddard, J. L., Paulsen, S. G. & Olsen, A. R. (In Press). Using relative risk to compare the effects of aquatic stressors at a regional scale. *Environmental Management*.

Nutrients

Bourassa, N., and A. Cattaneo. 1998. Control of periphyton biomass in Laurentian streams (Quebec). *J. N. Am. Benthol. Soc.*, 17:420–429.

Dodds, W.K. and E.B.Welch. 2000. Establishing nutrient criteria in streams. *J. N. Am. Benthol. Soc.*, 2000, 19(1):186–196

Kelly, M. G. 1998. Use of community-based indices to monitor eutrophication in rivers. *Environmental Conservation* 25:22–29.

Kelly, M. G., and B. A. Whitton. 1995. The trophic diatom index: a new index for monitoring eutrophication in rivers. *Journal of Applied Phycology* 7:433–444

Miltner, R. J., and E. T. Rankin. 1998. Primary nutrients and the biotic integrity of rivers and streams. *Freshwater Biology* 40:145–158.

Nordin, R. N. 1985. Water quality criteria for nutrients and algae (technical appendix). Water Quality Unit, Resources Quality Section, Water Management Branch, British Columbia Ministry of the Environment, Victoria, BC. (Available from: <http://www.env.gov.bc.ca> or Water Quality Section, BC Environment, PO Box 9340, Station Provincial Government, Victoria, BC V8W 9M1.)

Pan, Y., R. J. Stevenson, B. H. Hill, A. T. Herlihy, and G. B. Collins. 1996. Using diatoms as indicators of ecological conditions in lotic systems: a regional assessment. *J. N. Am. Benthol. Soc.* 15:481–495.

Welch, E. B. 1992. Ecological effects of wastewater. 2nd edition. Chapman and Hall, London, UK.